

Evidence for gravitational quadrupole moment variations in the companion of PSR J2051–0827

K. Lazaridis,¹ J. P. W. Verbiest,¹ T. M. Tauris,^{2,1} B. W. Stappers,^{3,4} M. Kramer,^{1,3} N. Wex,¹ A. Jessner,¹ I. Cognard,^{5,6} G. Desvignes,⁵ G. H. Janssen,³ M. B. Purver,³ G. Theureau,^{5,6} C. G. Bassa,³ R. Smits^{3,4}

¹Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

²Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

³University of Manchester, Jodrell Bank Centre for Astrophysics, Alan Turing Building, Manchester M13 9PL, UK

⁴Stichting ASTRON, Postbus 2, 7990 AA Dwingeloo, The Netherlands

⁵Laboratoire de Physique et Chimie de l’Environnement, CNRS, 3A Avenue de la Recherche Scientifique, 45071 Orléans Cedex 2, France

⁶Station de Radioastronomie de Nançay, Paris Observatory, France

Drafted 22 November 2010.

ABSTRACT

We have conducted radio timing observations of the eclipsing millisecond binary pulsar J2051–0827 with the European Pulsar Timing Array network of telescopes and the Parkes radio telescope, spanning over 13 years. The increased data span allows significant measurements of the orbital eccentricity, $e = (6.2 \pm 1.3) \times 10^{-5}$ and composite proper motion, $\mu_t = 7.3 \pm 0.4 \text{ mas yr}^{-1}$. Our timing observations have revealed secular variations of the projected semi-major axis of the pulsar orbit which are much more extreme than those previously published; and of the orbital period of the system. Investigation of the physical mechanisms producing such variations confirm that the variations of the semi-major axis are most probably caused by classical spin-orbit coupling in the binary system, while the variations in orbital period are most likely caused by tidal dissipation leading to changes in the gravitational quadrupole moment of the companion.

Key words: binaries: eclipsing - pulsars: general - pulsars: individual: PSR J2051–0827 - stars: evolution - stars: fundamental parameters

1 INTRODUCTION

PSR J2051–0827 is the second eclipsing millisecond pulsar system discovered after the original black widow pulsar PSR B1957+20 (Fruchter et al. 1988). It was discovered (Stappers et al. 1996) as part of a Parkes all-sky survey of the southern sky for low-luminosity and millisecond pulsars. The pulsar has a spin period of 4.5 ms and inhabits a very compact circular orbit with a very low mass companion: $m_c < 0.1 M_\odot$. The orbital period is $P_b = 2.38 \text{ h}$ and the pulsar and its companion are separated by just $1.0 R_\odot$. Stappers et al. (1996, 2001a) investigated the eclipse characteristics of this system and found that their duration is $\sim 10\%$ of the orbital period at frequencies below 1 GHz. In addition, assuming a mean free electron density of 10^7 cm^{-3} in the eclipse region, they calculated a mass-loss rate of $\sim 10^{-14} M_\odot \text{ yr}^{-1}$, which is insufficient to evaporate the companion completely within a Hubble time.

Optical observations of the companion revealed that it is irradiated by the pulsar wind (Stappers et al. 1996). Further observations and modelling of the variability of the light curve (Stappers et al. 1999; Stappers et al. 2001b) determined a binary inclination angle of $i \sim 40^\circ$, a backside temperature of $T \leq 3000 \text{ K}$,

a companion mass of $\sim 0.04 M_\odot$ and a radius $\sim 0.064 R_\odot$. An alternative model with a companion almost filling its Roche lobe ($\sim 0.12 R_\odot$) was also considered and produced almost the same results, albeit with a significantly worse fit. For clarification purposes, a cartoon of the system is given in Figure 1.

Doroshenko et al. (2001) presented the most precise timing analysis of PSR J2051–0827 using $\sim 6 \text{ yr}$ of radio timing measurements with the Effelsberg 100-m radio telescope and the 76-m Lovell at Jodrell Bank. Among the most important measurements is the variation of the projected semi-major axis, probably caused by Newtonian spin-orbit coupling in this binary system. In addition, they measured significant orbital period variations and concluded that those were created by the same mechanism as in the B1957+20 system (Applegate & Shaham 1994; Arzoumanian et al. 1994): by tidal dissipation of a tidally powered, non-degenerate companion.

Because black widow (and other eclipsing) systems are susceptible to variations of their orbital parameters, monitoring over long lengths of time is crucial to understand these systems. Timing of black widow pulsars has been undertaken by Nice et al. (2000) and Freire et al. (2003, 2005), but only very few of these

systems have been monitored for a decade or longer. This is of particular interest given the large number of black widow pulsars recently found in unidentified *Fermi* sources (Ransom et al. 2011; Keith et al. 2011, and several discoveries that are soon to be published).

In this paper we revisit PSR J2051–0827 with 13 years of high-precision timing data and combined datasets from the Australia Telescope National Facility’s Parkes 64-m radio telescope and the European Pulsar Timing Array (EPTA) telescopes, consisting of the 100-m Effelsberg radio-telescope of the Max-Planck-Institute for Radioastronomy, Germany, the 76-m Lovell radio-telescope at Jodrell Bank observatory of the University of Manchester, UK, the 94-m-equivalent Westerbork Synthesis Radio Telescope (WSRT) of ASTRON, the Netherlands, and the 94-m-equivalent Nançay decimetric Radio Telescope (NRT) of the CNRS, France. After describing the properties of our multi-telescope data (Section 2), we present the updated measurements of the astrometric, spin and binary parameters for the system (Section 3). Specifically we show the measurement of the combined proper motion and of the orbital eccentricity of the PSR J2051–0827 system for the first time and we measure the dispersion measure (DM) variations over time. Furthermore, we present the extreme variations in the orbital period and projected semi-major axis of the system, which are much larger than the ones published by Doroshenko et al. (2001). In Section 4 we rule out several possible contributions to the aforementioned variations and we discuss the physical mechanisms possibly responsible for those. Specifically, we show that gravitational quadrupole coupling (GQC) and classical spin-orbit coupling (SOC) can be viable mechanisms for the orbital variations of the PSR J2051–0827 system, under certain assumptions. In light of the new orbital variation measurements, we discuss the possible scenarios for the nature of the companion and the prospects for observations of the system at higher energies, also in Section 4. Finally, in Section 5, we briefly summarise our findings.

2 OBSERVATIONS AND DATA ANALYSIS

Description of the observing systems and the data acquisition procedure of the EPTA telescopes can be found in Lazaridis et al. (2009), while for Parkes this information is given in Stappers et al. (1998). A detailed description on combining pulse times-of-arrival (TOAs) from different telescopes can be found in Janssen et al. (2008); Lazaridis et al. (2009); details specific to the data analysis used here, are listed below.

In our timing of PSR J2051–0827, all TOAs with uncertainties greater than $20\mu\text{s}$ were excluded as they don’t impact the weighted fit. Furthermore, 573 TOAs that were taken at orbital phases between 0.2 and 0.35 were not used because the potential excess column density in the eclipsing region may cause additional modulation on the TOAs derived from these data (Stappers et al. 1998; Stappers et al. 2001a).

Timing was performed with the TEMPO2 software package (Hobbs et al. 2006), using the DE405 Solar-System ephemerides of the Jet Propulsion Laboratory (Standish 1998, 2004) and the ELL1 binary model (Lange et al. 2001). TEMPO2 minimises the weighted sum of the squared residuals, producing a set of improved pulsar parameters and post-fit timing residuals. In order to propagate unmodelled noise through into the uncertainties of the timing model parameters, the TOA uncertainties were multiplied by telescope-specific scaling factors to achieve a reduced χ^2 of unity.




Figure 2. Post-fit timing residuals for the last 4.5 years of data. Our timing solution is given in Table 2. The majority of the data are from the Nançay radio observatory, but data from Effelsberg, Jodrell Bank and Parkes were also used in the preceding nine years, which are not shown.

These scaling factors all fell between 1.4 and 3.5, where the largest factor was applied to the TOAs of the Effelsberg-Berkeley-Pulsar-Processor (EBPP). The large reduced χ^2 for the EBPP data is most likely caused by the fact that this instrument only records total power (no polarisation information) and has no RFI mitigation applied to it. Arbitrary phase offsets (also called “jumps”) were introduced between data obtained at different frequencies and/or taken with different telescopes, to account for frequency-dependent pulse shape evolution and observatory-dependent differences in instrumental delays, cable lengths and geodetic position, amongst others. These jumps were included as free parameters in the least-squares fit performed by TEMPO2 and they absorb any non-changing dispersive effects in our data. Therefore we were unable to evaluate the average DM and hence fixed the value obtained by Stappers et al. (1998) as a constant in our timing model. However, the data does contain some sensitivity to time-varying dispersive delays, which is fully discussed in Section 3.3.

In Table 1 the properties of the data from the different telescopes are presented. Combination of the TOAs from all the telescopes provides us with a 13-year dataset with no significant gaps. In order to obtain a coherent timing solution for this combined dataset, higher order derivatives (up to sixth order) of the orbital period and projected semi-major axis must be fitted, leading to high correlations between the modelled parameters. These variations in orbital period and size are discussed in detail in sections 4.1 and 4.2. To avoid these correlations and because 85% of our data (and all of our highest-precision data) were taken after 2004, we derive the timing model, presented in Table 2, based solely on the last 4.5 years of data, between MJDs 53293 and 54888 (as shown in Figure 2). The remaining older data were exclusively used to investigate potential variations in interstellar dispersion (see Section 3.3) and to investigate possible causes for the variations in orbital period and projected semi-major axis (Section 3.5).

Table 2. Timing parameters for PSR J2051–0827 for the last epoch of observations (MJD range 53293–54888).

Parameters	EPTA
Right ascension, α (J2000)	20 ^h 51 ^m 07 ^s .51808(2)
Declination, δ (J2000)	−08°27′37″.7608(9)
μ_α (mas yr ^{−1})	6.6(2)
μ_δ (mas yr ^{−1})	3.2(7)
ν (Hz)	221.796283737706(1)
$\dot{\nu}$ (s ^{−2})	−6.2639(9) × 10 ^{−16}
P (ms)	4.50864181828489(2)
$\dot{P}$ (s s ^{−1})	1.2733(2) × 10 ^{−20}
Reference epoch (MJD)	54091
MJD range of global timing model	53293 – 54888
MJD range epoch 1	49989 – 51545
MJD range epoch 2	51384 – 52332
MJD range epoch 3	51967 – 53117
MJD range epoch 4	52790 – 54248
MJD range epoch 5	54156 – 54888
Dispersion measure, DM (cm ^{−3} pc) *	20.7458(2)
Orbital period, P_b (days)	0.09911024846(2)
Projected semi-major axis, x (lt-s)	0.0450725(4)
η ($\equiv e \sin \omega$)	5(1) × 10 ^{−5}
κ ($\equiv e \cos \omega$)	4(1) × 10 ^{−5}
Eccentricity, e *	6(1) × 10 ^{−5}
Longitude of the periastron, ω (deg) *	52(12)
T_{ASC} (MJD)	54091.0343503(1)
$\dot{P}_b$, epoch 1 (s s ^{−1})	−1.33(6) × 10 ^{−11}
$\dot{P}_b$, epoch 2 (s s ^{−1})	0.9(1) × 10 ^{−11}
$\dot{P}_b$, epoch 3 (s s ^{−1})	1.8(3) × 10 ^{−11}
$\dot{P}_b$, epoch 4 (s s ^{−1})	−1.81(3) × 10 ^{−11}
$\dot{P}_b$, epoch 5 (s s ^{−1})	1.34(6) × 10 ^{−11}
$\dot{x}$, epoch 1 (s s ^{−1})	−1.7(4) × 10 ^{−13}
$\dot{x}$, epoch 2 (s s ^{−1})	−7.9(8) × 10 ^{−13}
$\dot{x}$, epoch 3 (s s ^{−1})	9(1) × 10 ^{−13}
$\dot{x}$, epoch 4 (s s ^{−1})	−0.8(2) × 10 ^{−13}
$\dot{x}$, epoch 5 (s s ^{−1})	0.2(2) × 10 ^{−13}
Solar system ephemeris model	DE405
Number of TOAs	3126
RMS timing residual (μ s)	12.2


* The eccentricity and the longitude of the periastron are calculated from the Laplace-Lagrange parameters, η and κ . The DM value and its uncertainty were taken from Stappers et al. (1998)

Figures in parentheses are the nominal 1 σ TEMPO2 uncertainties in the least-significant digits quoted.

These parameters were determined with TEMPO2, which uses the International Celestial Reference System and Barycentric Coordinate Time. Refer to Hobbs et al. (2006) for information on modifying this timing model for observing systems that use TEMPO format parameters.

3.3 DM variations

As described in Section 1, low-frequency observations of this system show eclipses as the pulsar passes through periastron. This may imply increased dispersion as a function of orbital phase, which could corrupt measurements of binary parameters. To investigate this possibility, we took the high-quality, multi-frequency data from the final 4.5 years and measured the DM as a function of orbital

**Figure 3.** DM variations versus time, including the best-fit linear trend.

phase while keeping the jumps between observing bands and observatories fixed. This did not result in significant trends at any orbital phase, including egress or ingress¹. Since this experiment required a fit for DM over TOAs that were restricted to a small fraction of the binary phase, a simultaneous fit to standard orbital parameters was impossible. However, the wide frequency range available at any given orbital phase implies that even if orbital-phase-dependent DM variations did occur, their effect on the TOAs could only partially be absorbed by the orbital parameters, implying some residual DM variations would still be visible. Simultaneous fitting for DM and long-term effects (such as pulsar position and proper motion, spin, spin-down and orbital derivatives $\dot{P}_b$ and $\dot{x}$) over limited orbital-phase ranges did not affect the DM estimates or the other timing parameters, confirming that the long-period terms of the timing model are not affected by any potential orbital-phase-dependent variations in interstellar dispersion.

In addition to investigating the DM evolution as a function of orbital phase, the DM variations were determined across the entire data set. To this end, intervals with adequate multi-frequency data were identified by hand and DM values were fitted in each interval, while jumps between telescopes and observing bands were kept fixed at the values determined from the entire data set. Clearly this cannot result in accurate measurements of DM, but it does allow variations to be determined precisely. The resulting measurements (shown in Fig. 3) demonstrate that no complex DM evolution is present in this data set, though a shallow DM decay is measured: $d(\text{DM})/dt = (-4.3 \pm 1.4) \times 10^{-4} \text{ cm}^{-3} \text{ yr}^{-1} \text{ pc}$. This measurement is barely significant at the 3- σ level. Given the small number of high-quality DM estimates that contributed to it and the possible underestimation of the DM measurement uncertainties (e.g. variations in orbital parameters could have affected the offset between non-simultaneous observations), we did not include a time-dependent DM variation in our timing model.

3.4 Companion mass

Optical observations (Stappers et al. 2001b) yielded a best fit for the orbital inclination angle of about 40°. Depending on the un-

¹ This does not contradict the increased column density in the eclipse region but underlines the limited sensitivity of our dataset to DM variations.




Figure 4. The masses of the stellar components in the J2051–0827 system as a function of orbital inclination angle. The box indicates the most likely parameter space in terms of the neutron star mass and constraints from radio eclipses and optical observations (see text).

known mass of the pulsar this results in a companion mass in the range: $m_c \simeq 0.04 - 0.06 M_\odot$ (see Figure 4) given the constraints from the mass function:

$$f = \frac{(m_c \sin i)^3}{(m_p + m_c)^2} = \frac{4\pi^2}{G} \frac{(a_p \sin i)^3}{P_b^2} = 1.0030 \times 10^{-5} M_\odot, \quad (2)$$

where the measured observables are the orbital period, P_b and the projected semi-major axis of the pulsar orbit, $a_p \sin i$. For the rest of this paper we shall assume a companion star mass of $m_c = 0.05 M_\odot$ and consequently a pulsar mass of $m_p = 1.8 M_\odot$. The pulsar mass of PSR J2051–0827 is likely to be significantly larger than the typical neutron star mass ($1.35 M_\odot$) obtained from measurements in double neutron star binaries, given that this system has evolved through a long ($\sim$ Gyr) LMXB phase with sub-Eddington mass transfer (see, e.g. Pfahl et al. 2002). Recent work on the mass determination of the original black widow pulsar (PSR B1957+20) by van Kerkwijk et al. (2011) confirms that the pulsars in these systems can accrete significant amounts of matter. For a discussion of the effect of irradiation on the accretion efficiency in LMXBs, see Ritter (2008). The Roche-lobe radius of the companion star in PSR J2051–0827 changes slightly with the estimated stellar masses and is found to be $R_L = 0.15 \pm 0.01 R_\odot$. However, the size (radius) of the irradiated companion star is difficult to determine accurately and in the discussion further on we shall assume two different values for the filling factor (the ratio of the volume-equivalent radius of the companion star to its Roche lobe) of 0.43 and 0.95 (Stappers et al. 2001b), corresponding to stellar radii of about $0.064 R_\odot$ and $0.14 R_\odot$ respectively.

3.5 Orbital changes

In order to monitor their variations over time, values for P_b and x were derived for each year of data, where three months of overlap were kept between adjacent years. In doing so, all model parameters besides x , P_b and T_{ASC} were held fixed and the timing reference epoch was defined to be the centre of each year-long interval. The fractional changes of these measurements are shown in Figure 5. Clearly, five different epochs can be identified, in which variations of both P_b and x can be described using only a linear trend, as shown in the figure.




Figure 5. Changes of the orbital period (top) and projected semi-major axis (bottom) versus date. Each point corresponds to close to one year of data, with typically three months overlap between adjacent points, though the precise amount of overlap varies depending on the density of observations and the amplitude of the orbital variations. The horizontal bars indicate the intervals over which measurements were made, and the uncertainties of the measurements are shown by vertical error bars that are placed at the mean TOA of the data contained in the interval. Because of inhomogeneous sampling, the mean TOA is not necessarily at the middle of the interval. For each of the five epochs listed in Table 2 a trend line derived from the respective $\dot{P}_b$ and $\dot{x}$ measurements given in Table 2 is shown. In most cases these trend lines fit well through the measurement points, though deviations may be caused by rapid variations in orbital parameters, underestimation of measurement uncertainties, correlations between parameters and the fact that on many of the shorter measurement intervals only part of the timing model could be fitted for, incurring potential corruptions in correlated parameters. The thick horizontal bars on the top of the figure indicate the timing baseline of Doroshenko et al. (2001), which only really spans the first two of these epochs.

4 DISCUSSION

4.1 Orbital period variations

$\dot{P}_b$ is the observable rate of change of the orbital period and is caused by a variety of effects, both intrinsic to the system and

caused by kinematic effects relative to the observer. The most important contributions are:

$$\dot{P}_b = \dot{P}_b^{\text{GW}} + \dot{P}_b^{\text{D}} + \dot{P}_b^{\text{m}} + \dot{P}_b^{\text{T}} + \dot{P}_b^{\text{Q}}. \quad (3)$$

As a point of reference, the values for $\dot{P}_b$ in the two most extreme epochs are $\dot{P}_b = -1.81(3) \times 10^{-11}$ and $\dot{P}_b = 1.8(3) \times 10^{-11}$.

The first term, $\dot{P}_b^{\text{GW}}$, is the contribution due to gravitational wave emission. In general relativity, for circular orbits it is given by (Peters 1964):

$$\dot{P}_b^{\text{GW}} = -\frac{192\pi}{5} \left(\frac{2\pi}{P_b} \frac{Gm_c}{c^3} \right)^{5/3} \frac{q}{(q+1)^{1/3}}, \quad (4)$$

The mass ratio $q = 36$ has been calculated assuming a pulsar mass $m_p = 1.8 M_\odot$ and a companion mass $m_c = 0.05 M_\odot$ for an inclination angle of $i = 40^\circ$. For PSR J2051–0827 we find $\dot{P}_b^{\text{GW}} \simeq -7.5 \times 10^{-14}$. This value is about three orders of magnitude less than the observed value of $\dot{P}_b$.

The second term, $\dot{P}_b^{\text{D}}$, is the Doppler correction, which is the combined effect of the proper motion of the system (Shklovskii 1970) and a correction term for the Galactic acceleration. The contribution for the Galactic acceleration at the location of PSR J2051–0827, $\dot{P}_b^{\text{Gal}}$, is of order 1.1×10^{-15} (Lazaridis et al. 2009). Using numbers from Table 2, we also calculate the contribution due to the Shklovskii effect according to the following:

$$\dot{P}_b^{\text{Shk}} = \frac{(\mu_\alpha^2 + \mu_\delta^2)d}{c} P_b \simeq 1.1 \times 10^{-15}. \quad (5)$$

By summing we yield the Doppler correction:

$$\dot{P}_b^{\text{D}} = \dot{P}_b^{\text{Gal}} + \dot{P}_b^{\text{Shk}} \simeq 2.2 \times 10^{-15}, \quad (6)$$

four orders of magnitude smaller than the measured value.

An acceleration of the binary system with respect to the Solar System Barycentre (SSB) could also be caused by a third massive body orbiting the binary system. However, it would affect the orbital period derivative and the spin period derivative in the same way. Assuming that the spin period derivative is caused entirely by the acceleration, we can estimate the maximal effect this would have on $\dot{P}_b$: $(\dot{P}_b/P_b)^{\text{acc}} = (\dot{P}/P) \simeq -3 \times 10^{-18} \text{ s}^{-1}$. This is seven orders of magnitude smaller than the measured value, thus there can be no massive third body orbiting the system.

The third term, $\dot{P}_b^{\text{m}}$, is the contribution from the mass loss of the binary. If we consider a circular orbit, no mass loss from the pulsar ($\dot{m}_p = 0$) and a rate of mass loss from the companion $\dot{m}_c > 0$ (Jeans 1924), we obtain:

$$\frac{\dot{P}_b}{P_b} = -2 \frac{\dot{m}_c}{M} \quad (7)$$

and therefore:

$$\dot{m}_c = -\frac{M}{2} \frac{\dot{P}_b}{P_b} \simeq 6.2 \times 10^{-8} M_\odot \text{ yr}^{-1}, \quad (8)$$

where M is the total mass of the system. This number is six orders of magnitude larger than expected (as calculated by Stappers et al. 1996) and hard to reconcile with the low electron densities measured in the eclipse region (Stappers et al. 1996). In addition, for the second of the epochs we would need mass injection to the companion in order to explain the orbital decay – an infeasible scenario for this system.

Tidal forces in tight synchronous binaries may also be responsible for interactions which would change the orbital period. This

could be the case if, for example, a companion star suddenly contracts or expands and thereby changes its spin angular momentum. Tidal locking of the orbit would then result in exchange of spin and orbital angular momentum, e.g. Tauris & Savonije (2001). However, in PSR J2051–0827 we do not see reasons for any sudden major radial changes in the structure of the companion star – especially not changes that would not result in detection of X-ray bursts (A. Levine and D. Altamirano, private communication). Hence, we neglect the term $\dot{P}_b^{\text{T}}$.

Since all the above contributions are much smaller than the observed variation of the orbital period, we conclude that they must originate from the last term of Equation (3), which represents variations of the gravitational quadrupole moment of the companion star.

4.1.1 The Applegate GQC model

The gravitational quadrupole coupling (GQC) mechanism (Applegate 1992; Applegate & Shaham 1994) has been applied successfully to the eclipsing binary system PSR B1957+20 (Applegate & Shaham 1994), explaining the orbital period variations seen by Arzoumanian et al. (1994). In addition, it was proposed by Doroshenko et al. (2001) as the main mechanism for producing the orbital period variations of PSR J2051–0827. However, as we will present below, the latter publication considered only a small part of these variations, which led to an underestimation in their calculations.

In the GQC model (Applegate & Shaham 1994), magnetic activity is driven by energy flows in convective layers of the irradiated companion. This, in combination with wind mass loss, results in a torque on its spin, which holds it slightly out of synchronous rotation, causing tidal dissipation of energy and heating of the companion. The resultant time-dependent gravitational quadrupole moment (e.g. variations of the oblateness) causes modulation of the orbital period on a short, in principle dynamical, time scale. An increase in the quadrupole moment causes the two stars to move closer together and a decrease in the quadrupole moment results in a widening of the orbit – in Applegate’s model total orbital angular momentum is assumed to remain constant and the neutron star is treated as a point mass. The details of the hydrodynamic dynamo and its activity cycle remain unspecified for our purposes (also given the unknown nature of the companion star – see Section 4.4) and it is simply assumed that the variable quadrupole moment, ΔQ , is caused by cyclic spin-up and spin-down of the outer layers of the companion, which leads to orbital period changes equal to (Applegate & Shaham 1994):

$$\left(\frac{\Delta P_b}{P_b} \right)^Q = -9 \frac{\Delta Q}{m_c a^2}, \quad (9)$$

where a is the separation and m_c the companion mass. The transfer of angular momentum ΔJ to a thin shell of radius R_c and mass M_s , rotating with angular velocity Ω in the gravitational field of a point mass m_c , will cause the quadrupole moment of the shell to change by (Applegate & Shaham 1994):

$$\Delta Q = \frac{2}{9} \frac{M_s R_c^5}{G m_c} \Omega \Delta \Omega, \quad (10)$$

where G is the Newtonian gravitational constant and $\Delta \Omega$ is the change in the angular velocity of the shell. In order to produce orbital period changes ΔP_b you need a variable angular velocity given by

$$\frac{M_s}{m_c} \frac{\Delta\Omega}{\Omega} = \frac{Gm_c}{2R_c^3} \left(\frac{a}{R_c}\right)^2 \left(\frac{P_b}{2\pi}\right)^2 \frac{\Delta P_b}{P_b} \quad (11)$$

(Applegate & Shaham 1994). This procedure produces a luminosity variation ΔL given by

$$\Delta L = \frac{\pi}{3} \frac{Gm_c^2}{R_c P_{\text{mod}}} \left(\frac{a}{R_c}\right)^2 \frac{\Omega_{\text{dr}}}{\Omega} \frac{\Delta P_b}{P_b}, \quad (12)$$

where P_{mod} is the period of the orbital period modulation and Ω_{dr} is the angular velocity of the differential rotation (Applegate & Shaham 1994).

Applegate & Shaham (1994) assume in their model a mass $M_s \simeq 0.1m_c$ for the thin shell; an angular velocity $\Omega_{\text{dr}} \simeq \Delta\Omega$; and luminosity variations at the $\Delta L/L \simeq 0.1$ level. From Figure 5 we calculated for PSR J2051–0827 a total change in the orbital period of $\Delta P_b/P_b \simeq 2.2 \times 10^{-7}$ and a modulation period of $P_{\text{mod}} \simeq 7.5$ yr. As mentioned above these two values were under- and over-estimated, respectively, in Doroshenko et al. (2001) because of their significantly smaller data set. By assuming an inclination angle of $i = 40^\circ$ and a companion radius of $R_c \sim 0.064 R_\odot$ (Stappers et al. 2001b) for a separation $a \simeq 7.1 \times 10^8$ m and a companion mass $m_c = 0.05 M_\odot$, we derive a variation of the angular velocity of the companion of:

$$\frac{\Delta\Omega}{\Omega} \sim 4 \times 10^{-2}. \quad (13)$$

This variation would produce a variable luminosity of $\Delta L \sim 1.4 \times 10^{33} \text{ erg s}^{-1}$ and according to the model the internal luminosity of the companion must be

$$L \sim 1.4 \times 10^{34} \text{ erg s}^{-1}. \quad (14)$$

From the optical observations of the companion (Stappers et al. 2001b) we can estimate a maximum effective temperature of $T_{\text{eff}}^{\text{max}} \simeq 3000$ K which gives an internal luminosity of $L \sim 10^{30} \text{ erg s}^{-1}$. We therefore conclude that, under the assumptions listed above, the GQC model produces much larger orbital period variations than expected and cannot fit the PSR J2051–0827 system. If the companion was tidally powered, the internal luminosity derived from the optical observations would be much higher. One reason why GQC does not fit the observations may be the simplifications of the model itself, i.e. the consideration that all of the energy variations appear as luminosity variations without any loss (Applegate 1992). Although improvement of the model could possibly decrease ΔL by a factor of a few, the difference we calculate is significantly larger.

The previous calculations can change by a large factor if we consider the alternative model that Stappers et al. (2001b) presented for the companion star, where the latter almost fills its Roche lobe ($R_c = 0.14 R_\odot$). Using the values for that model, we derive a variation of the angular velocity of the companion of

$$\frac{\Delta\Omega}{\Omega} \sim 7.1 \times 10^{-4} \quad (15)$$

and an internal luminosity of

$$L \sim 2 \times 10^{31} \text{ erg s}^{-1}. \quad (16)$$

This number is much closer to the newly calculated internal luminosity from the optical of $L \sim 6 \times 10^{30} \text{ erg s}^{-1}$. Thus, under specific assumptions the GQC model can explain the orbital period variations of PSR J2051–0827.

4.2 Changes in the projected semi-major axis

The observed values of $\dot{x}$ can be the result of various effects (Lorimer & Kramer 2005):

$$\dot{x}_{\text{obs}} = \dot{x}^{\text{GW}} + \dot{x}^{\text{D}} + \frac{d\epsilon_A}{dt} + \dot{x}^{\text{PM}} + \dot{x}^{\text{m}} + \dot{x}^{\text{Q}} + \dot{x}^{\text{SOC}}. \quad (17)$$

For the most extreme epochs, we have: $\dot{x} = -7.9(8) \times 10^{-13}$ and $\dot{x} = 9(1) \times 10^{-13}$.

The first term, $\dot{x}^{\text{GW}}$ arises from orbital shrinkage due to gravitational-wave damping. Using Kepler’s third law and Equation 4:

$$\frac{\dot{x}^{\text{GW}}}{x} = \frac{2}{3} \frac{\dot{P}_b^{\text{GW}}}{P_b} \simeq -2.6 \times 10^{-19} \quad (18)$$

(Peters 1964). This contribution is much smaller than the current measurement.

The second term, $\dot{x}^{\text{D}}$, is identical to the second term of Equation (3). The contribution for the Galactic acceleration is of order 1.1×10^{-15} and the contribution of the Shklovskii effect $\dot{x}^{\text{Shk}} = x(\mu_\alpha^2 + \mu_\delta^2)d/c \sim 6.0 \times 10^{-21}$. Both of these are very small compared to the observed value so this term can be neglected.

The third term, $d\epsilon_A/dt$, is the contribution of the varying aberration caused by geodetic precession of the pulsar spin axis and is typically of order $\Omega^{\text{geod}} P/P_b \approx 7.4 \times 10^{-17}$ (Damour & Taylor 1992). For a recycled pulsar, like PSR J2051–0827, the spin is expected to be close to parallel to the orbital angular momentum, which further suppresses this effect. Hence, the contribution is at least three orders of magnitude smaller than the observed value.

The fourth term, $\dot{x}^{\text{PM}}$, represents a variation of x caused by a change of the orbital inclination while the binary system is moving relatively to the SSB (Arzoumanian et al. 1996; Kopeikin 1996). This effect is quantified by the following equation:

$$\dot{x}^{\text{PM}} = 1.54 \times 10^{-16} x \cot i (-\mu_\alpha \sin \Omega_{\text{asc}} + \mu_\delta \cos \Omega_{\text{asc}}), \quad (19)$$

where Ω_{asc} is the position angle of the ascending node. The quantities x , μ_α and μ_δ are expressed in seconds and milliarcseconds per year, respectively. The maximal contribution of the proper motion is:

$$\dot{x}_{\text{max}}^{\text{PM}} = 1.54 \times 10^{-16} x (\mu_\alpha^2 + \mu_\delta^2)^{1/2} \cot i. \quad (20)$$

For an inclination angle of $i = 40^\circ$, we get $\dot{x}_{\text{max}}^{\text{PM}} \simeq 6.0 \times 10^{-17}$. Thus, this term is also very small compared to the measured $\dot{x}$.

The fifth term, $\dot{x}^{\text{m}}$, represents a change in the size of the orbit caused by mass loss from the binary system. Using Equation (7) and Kepler’s third law, we calculate the rate of mass loss from the companion to be seven to eight orders of magnitude larger than expected.

The sixth term, $\dot{x}^{\text{Q}}$, is caused directly by the change in quadrupole moment as described in Section 4.1. Because orbital angular momentum is conserved in the Applegate model, a change in the orbital period is related to a change in the size of the orbit by

$$\frac{\Delta a}{a} = 2 \frac{\Delta P_b}{P_b}. \quad (21)$$

It can be seen from Figure 5 that this contribution falls short by four orders of magnitude to explain the observed changes in the projected semi-major axis x .

As all other contributions are much smaller than the observed variation of the projected semi-major axis, they must originate from the last term of Equation (17): the classical spin-orbit coupling (SOC) term.

Table 3. Quadrupole moment variations for the five epochs, as derived from the P_b variations, alongside the x variations as observed.

Epoch	ΔQ (10^{45} g cm^2)	$n_b \Delta \tilde{Q}$ (10^{-11} s^{-1})	$\dot{x}/x$ (10^{-11} s^{-1})
1	11.2	7.7	-0.38
2	-4.6	-3.2	-1.8
3	-11.3	-7.7	2.0
4	14.3	9.8	-0.18
5	-5.3	-3.6	0.044

4.2.1 The Applegate GQC model with SOC

Through SOC, the quadrupole of a rapidly rotating companion leads to apsidal motion and precession of the binary orbit. This in turn causes a variation of the longitude of periastron (which is impossible to measure in a system with such small eccentricity) and of the projected semi-major axis, according to:

$$\dot{x}^{\text{SOC}} = x n_b \tilde{Q} \cot i \sin \theta \cos \theta \sin \Phi, \quad (22)$$

(Smarr & Blandford 1976; Wex 1998), where $n_b = 2\pi/P_b$ is the orbital frequency, θ is the angle between the spin and orbital angular momentum and Φ is the longitude of the ascending node with respect to the invariable plane (plane perpendicular to the total angular momentum). The dimensionless quadrupole $\tilde{Q}$ is related to the quadrupole Q by

$$\tilde{Q} = \frac{3}{2} J_2 \left(\frac{R_c}{a} \right)^2, \quad J_2 = \frac{3Q}{m_c R_c^2}, \quad (23)$$

with J_2 the dimensionless measure of the quadrupole moment. Since no uniform change in x is detected, we can consider the companion to consist of a long-term stable component with spin axis aligned to the orbital angular momentum axis (i.e. $\theta \approx 0$) and in addition to that, a part of the star (like e.g. an outer shell) that changes its quadrupole moment by $\Delta \tilde{Q}$ and has an effective and variable angle θ independent of the rest of the star. This part will then give rise to a variation $\dot{x}/x$ of the order $n_b \Delta \tilde{Q}$.


In Section 4.1.1 we have described how changes in the gravitational quadrupole moment of the companion could cause the observed P_b variations. Even though these direct GQC effects are insufficient to explain the x variations (i.e. $\dot{x}^{\text{Q}} = 0$ as explained above), through the SOC mechanism they might have a more substantial effect.

Extending Equation 9 to arbitrary θ and using Equation 23, we have:

$$\frac{\Delta P_b}{P_b} = -2\Delta \tilde{Q} \left(1 - \frac{3}{2} \sin^2 \theta \right). \quad (24)$$

Combining this with the observed $\dot{P}_b$ values (Table 2) and assuming $\theta = 0$ for now, we get the maximum $\Delta \tilde{Q}$ values presented in Table 3. Comparison to the $\dot{x}/x$ values from Figure 5, also listed in Table 3, shows that while the order of magnitude is roughly correct, the sign does neither correlate nor anti-correlate, which means that the orientation of the quadrupole-moment changes (i.e. θ) must vary strongly from epoch to epoch, which may be unphysical.

In order to examine more closely if the orbital variations can be produced by GQC and SOC arising from the same ΔQ , we have plotted in Figure 6 how $|\Delta \tilde{Q}|$ depends on θ , assuming $\Phi = 45^\circ$ and using $i = 40^\circ$. The curves are plotted for SOC, as derived from Equation 22, and for GQC, as derived from Equation 24. The intersection of the two lines clearly shows that for several values of

**Figure 6.** Quadrupole moment changes versus θ as produced by SOC (solid line) and GQC (dashed line) for epochs two (top) and three (bottom).

θ a $\Delta \tilde{Q}$ could explain both the orbital period and projected semi-major axis variations for PSR J2051–0827, though these θ values change significantly with epoch and we do not see how this easily fits with the physical mechanism behind the thin shell model of Applegate (1992) (see also our Section 4.1.1).

4.2.2 Spin precession of the companion star

In Section 4.2.1 we assumed a mostly stable companion star with an outer shell responsible for the changing quadrupole moment. An alternative explanation for the variations in P_b and x could be provided by limited quadrupole changes (as needed for the P_b variations, see Section 4.1.1) combined with an overall small tilt θ of the entire companion star with respect to the orbit. This tilt causes a precession of both the star and the orbit and it allows the entire quadrupole moment of the star $\tilde{Q}$ to cause variations in x according to Equation 22. We stress that in this scenario there need not be a physical mechanism relating the spin precession (which causes the

Table 4. Quadrupole changes (orientation and magnitude) that explain the changes in P_b and x for the five epochs.

Epoch	θ (deg)	$\Delta Q/Q$
1	−0.14	+0.041
2	−0.64	−0.017
3	+0.73	−0.041
4	−0.06	+0.052
5	+0.02	−0.019

x variations) to the quadrupole moment changes (that induce the P_b changes)².

To evaluate this model, we estimate the quadrupole moment of the companion star assuming a non-degenerate structure³ and following Lanza & Rodonò (1999):

$$Q = \frac{5}{6\xi} \frac{R^3 [\mathcal{T} - (1 - 3\eta)\mathcal{M}]}{GM} \quad (25)$$

where R and M are the radius and mass of the star, $\mathcal{T}$ is its rotational kinetic energy ($\mathcal{T} = I\Omega^2/2$), $\mathcal{M}$ is its total magnetic energy, η is the fraction of this magnetic energy in a field directed along the rotation axis and ξ is a dimensionless parameter depending on the density stratification and the spatial distribution of the perturbing forces in the convective layer of the star (it can be calculated from the gyration radius and the apsidal motion constant using stellar models). Here we shall use $\xi \simeq 5$. The rotational frequency of the outer layers of the star is taken to be that of the orbit (caused by synchronisation) and in Applegate’s model $\mathcal{M} = 0$. Adopting a companion star radius $R = 0.064 R_\odot$ yields:

$$Q = 3.4 \times 10^{47} \text{ g cm}^2.$$

Consequently, the quadrupole moment Q of the star is one to two orders of magnitude larger than the variations required for the observed P_b changes. Using this value in Equation 22 results in the angles θ required for the observed x variations. These values can in turn be used to correct the ΔQ values shown in Table 3 for the angular dependence, though this effect is all but negligible. The resulting changes required in the tilt and quadrupole moment of the companion star are collated in Table 4.

We conclude that with small changes in the orientation and magnitude of the quadrupole moment, we can explain all observed parameter changes. The change in the sign of $\dot{x}$ from one epoch to the other can be easily explained by a small oscillation of the symmetry axis of the quadrupole. Moreover, if we use an alternative scenario with a larger filling factor for the radius of the companion star, these numbers will get even smaller since Q will be larger. We do not, however, know of any physical mechanism that would cause the required tilt of the star or its rapid changes.

² Note the changes in θ will have some contribution to the P_b changes as well. Based on Equation 5 of Applegate (1992) we determined this effect and for all epochs save the third, this effect is at or well below the few percent level of the P_b changes induced by ΔQ . In the third epoch, which requires the largest angle θ , it contributes 1% of the observed P_b variations.

³ Deviations of a semi-degenerate nature should not be important in this case.

4.3 Alternative Models for the Gravitational Quadrupole-Moment Variations

The Applegate model is one of a number that have been applied to explain the cyclic modulation in the orbital period of magnetically active close binaries. In Applegate’s model a rather large fraction of the stellar luminosity is required for its operation. Hence, one might be able to detect changes in the stellar luminosity in phase with the orbital period modulation. The model of Lanza & Rodonò (e.g. Lanza et al. 1998; Lanza & Rodonò 1999) considers variations in the azimuthal B-field to explain the variations in oblateness. A change in the azimuthal field intensity can also produce a change in the quadrupole moment by changing the effective centrifugal acceleration. However, one should keep in mind that it remains to be established what kind of companion star we have in PSR J2051–0827.

4.4 The Nature of the Companion Star

There seem to be three possibilities for the nature of the companion star:

- I a white dwarf (WD),
- II a brown-dwarf-like star and
- III a semi-degenerate helium star (He).

In the following we briefly discuss each of these possibilities.

For a WD companion star of $M_c \geq 0.04 M_\odot$, the mass-radius relation of a non-relativistic degenerate Fermi-gas (e.g. Shapiro & Teukolsky 1983) yields an upper limit for the radius of $R_c \leq 0.025 R_\odot$. This value is much less than what is estimated in optical since a combination of the minimum value for the filling factor of 0.43 and the Roche-lobe size of $R_L = 0.15 R_\odot$, yields a minimum optical radius of $R_c \geq 0.064 R_\odot$. However, because of irradiation by the pulsar wind and tidal dissipation of energy in the WD envelope it is likely that the WD is bloated in size. The effect of a thermally bloated WD in a close binary system has recently been detected in a transiting source by the Kepler satellite (Carter et al. 2011). In this case the young (few hundred Myr) hot white dwarf is bloated by a factor of seven in size. The companion star in PSR J2051–0827 is believed to be much older, however it is quite possible that an extended H-rich atmosphere may exist in this star from the effects mentioned above. On the other hand, the actual effect of the pulsar wind remains uncertain and may not be as efficient as expected previously. For example, Equation (2) of Tavani (1992) for the irradiation of the companion as induced by the pulsar wind, yields a surface temperature of the companion of 6200 K. However, the observed temperature is ≤ 3000 K (Stappers et al. 2001b).

Brown dwarf models have been applied (Bildsten & Chakrabarty 2001) to explain the nature of the $\sim 0.05 M_\odot$ donor star in the accreting millisecond pulsar system SAX J1808.4–3658. Brown dwarfs have a relation between mass and radius that is acceptable from observational constraints, also for PSR J2051–0827. However, the obtained mass-radius relation was based on brown-dwarf models by Chabrier et al. (2000), calculated for isolated low-mass stars. We find it questionable if the present core remnant in PSR J2051–0827 can simply be described by the equivalent of a brown dwarf. The original ZAMS progenitor system of PSR J2051–0827 definitely did not consist of a B-star and a brown dwarf: the initial sub-stellar mass of a brown dwarf is $< 0.08 M_\odot$ (the limit for H-ignition) and the progenitor of the neutron star had a mass $> 10 M_\odot$. Hence, the

initial binary would have had a mass ratio of $< 1/100$, which is not only disfavoured by formation, but also unlikely given that this binary should undergo a common envelope phase followed by survival of a supernova explosion and finally spin up the pulsar with so little mass. A more reasonable possibility is that PSR J2051–0827 (and many other eclipsing binary millisecond pulsars) is the outcome of a short orbital period, low-mass X-ray binary with un-evolved main sequence stars. Such systems have been studied in detail by, e.g., Pylyser & Savonije (1988) and Ergma et al. (1997). The latter authors also discussed the final fate of a system like PSR J2051–0827: an ultra-compact LMXB where the companion fills its Roche lobe (again), because of orbital angular momentum losses, and eventually may undergo tidal disruption leaving behind a single or planetary millisecond pulsar (e.g. a system like PSR B1257+12, Wolszczan & Frail 1992).

Finally, it should be noted that the companion star in PSR J2051–0827 could be closely related to the semi-degenerate helium star companions in ultra-compact AM CVn systems (e.g. Iben & Tutukov 1991; Nelemans et al. 2001). One example is the system SDSS J0926+3624 (Copperwheat et al. 2010) which has a similar companion mass of $0.035 M_{\odot}$ ($P_b = 28$ min). Although the accreting compact object in AM CVn systems is a WD, a similar evolution is expected for systems with an accreting neutron star (Savonije et al. 1986). Also, in this case the expected mass-radius relations yield acceptable values for the radius in PSR J2051–0827 – especially if one accepts the possibility that a pulsar wind and tidal dissipation could lead to a somewhat bloated radius.

4.5 High-Energy Emission Prospects for PSR J2051–0827

The detection of X-rays from PSR J2051–0827 could yield vital information about its orbital evolution, the state of the companion star and interactions between the pulsar wind and the evaporated material of the companion star. However, no X-rays are detected from PSR J2051–0827 in the RXTE All-Sky-Survey (A.M. Levine, private communication). The RXTE ASM sensitivity in the 2–10 keV band is ~ 30 mCrab, corresponding to an energy flux of $F_x \simeq 7 \times 10^{-10} \text{ erg s}^{-1} \text{ cm}^{-2}$. Given the distance to PSR J2051–0827 ($d \simeq 1.04$ kpc), we can estimate an upper limit to the mass accretion rate:

$$\dot{m} \leq \frac{4\pi d^2 F_x R_p}{G m_p} \quad (26)$$

Assuming a neutron star mass $m_p = 1.8 M_{\odot}$ and radius $R_p = 10^6$ cm, we obtain a conservative upper limit for the potential accretion rate $\dot{m} \lesssim 6.0 \times 10^{-12} M_{\odot} \text{ yr}^{-1}$. Keeping in mind that the expected mass transfer rate for a Roche-lobe-filling $\sim 0.05 M_{\odot}$ companion star is expected to be much lower than this limit – and given the fact that we do observe a radio pulsar – it is not a surprise that PSR J2051–0827 is not detected in the RXTE ASM data. Chandra and/or XMM data would impose far more stringent constraints; or might even provide a detection.

One must bear in mind that the lack of detected X-rays does not exclude mass loss from the companion star. It is possible that the pulsar wind (which is probably enhanced in the equatorial region towards the companion star) is able to prevent any accretion onto the neutron star. However, in this scenario it needs to be investigated thoroughly whether or not a shock front would lead to acceleration of protons and subsequent production of high-energy γ -rays (Harding & Gaisser 1990; Stappers et al. 2003).

Considering γ -ray emission from this system, with a distance of $\simeq 1.04$ kpc and a spin-down luminosity $\dot{E} = 4\pi^2 \dot{I} \dot{P} / P^3 =$

$5.5 \times 10^{33} \text{ erg/s}$, this pulsar is a good candidate for detection with the Large Area Telescope of the *Fermi* satellite (Abdo et al. 2009). However none of the *Fermi* sources in the first-year catalogue lie within a radius of 3° of PSR J2051–0827 (Abdo et al. 2010). It would therefore be interesting to see if PSR J2051–0827 is detected by *Fermi* with accumulated data.

5 SUMMARY

We have presented a timing update on PSR J2051–0827 and shown that the variations of the orbital period and projected semi-major axis of the binary system are far more extreme than described in earlier work. We have analysed all possible causes for these variations and found that a combination of gravitational quadrupole coupling and spin-orbit coupling are the most likely origins of this behaviour, though accurate modelling efforts are still required. We furthermore discussed the nature of the companion star and conclude that a semi-degenerate helium star is more likely than the white-dwarf alternative and, finally, we expect high-energy γ -rays may be visible in this system if a pulsar wind prevents accretion onto the neutron star – which in turn is expected based on the lack of X-rays.

ACKNOWLEDGEMENTS

We are very grateful to all staff at the Effelsberg, Westerbork, Jodrell Bank and Nançay radio telescopes for their help with the observations used in this work. Part of this work is based on observations with the 100-m telescope of the Max-Planck-Institut für Radioastronomie (MPIfR) at Effelsberg. Access to the Lovell telescope is supported through an STFC rolling grant. The Nançay radio telescope is part of the Paris Observatory, associated with the Centre National de la Recherche Scientifique (CNRS) and partially supported by the Région Centre in France. The Westerbork Synthesis Radio Telescope is operated by the Netherlands Foundation for Research in Astronomy (ASTRON) with support from the NWO. The Parkes radio telescope is part of the Australia Telescope National Facility which is funded by the Commonwealth of Australia for operation as a national facility managed by CSIRO. JPWV is supported by the European Union under Marie-Curie Intra-European Fellowship 236394. We want to thank Alan M. Levine (MIT) and Diego Altamirano (University of Amsterdam) for their help with investigating RXTE data, Lucas Guillemot (MPIfR) for discussions on Fermi results and Paulo Freire (MPIfR) for discussions on black widow and eclipsing systems. Finally, we thank the referee, Ingrid Stairs (UBC), for helpful and interesting comments.

REFERENCES

- Abdo A. A., et al., 2009, *Science*, 325, 848
- Abdo A. A., et al., 2010, *ApJS*, 18, 405
- Applegate J. H., 1992, *ApJ*, 385, 621
- Applegate J. H., Shaham J., 1994, *ApJ*, 436, 312
- Arzoumanian Z., Fruchter A. S., Taylor J. H., 1994, *ApJ*, 426, L85
- Arzoumanian Z., Joshi K., Rasio F., Thorsett S. E., 1996, in Johnston S., Walker M. A., Bailes M., eds, *Pulsars: Problems and Progress*, IAU Colloquium 160 Orbital parameters of the PSR B1620–26 triple system. Astronomical Society of the Pacific, San Francisco, pp 525–530
- Bildsten L., Chakrabarty D., 2001, *ApJ*, 557, 292

- Carter J. A., Rappaport S., Fabrycky D., 2011, *ApJ*, 728, 139
- Chabrier G., Baraffe I., Allard F., Hauschildt P., 2000, *ApJ*, 542, 464
- Copperwheat C. M., Marsh T. R., Littlefair S. P., Dhillon V. S., Ramsay G., Drake A. J., Gänsicke B. T., Groot P. J., Hakala P., Koester D., Nelemans G., Roelofs G., Southworth J., Steeghs D., Tulloch S., 2010, *MNRAS*, pp 1557–+
- Cordes J. M., Lazio T. J. W., 2002
- Damour T., Taylor J. H., 1992, *Phys. Rev. D*, 45, 1840
- Doroshenko O., Löhmer O., Kramer M., Jessner A., Wielebinski R., Lyne A. G., Lange C., 2001, *A&A*, 379, 579
- Ergma E., Lundgren S. C., Cordes J. M., 1997, *ApJ*, 475, L29
- Freire P. C., Camilo F., Kramer M., Lorimer D. R., Lyne A. G., Manchester R. N., D’Amico N., 2003, *MNRAS*, 340, 1359
- Freire P. C. C., Hessels J. W. T., Nice D. J., Ransom S. M., Lorimer D. R., Stairs I. H., 2005, *ApJ*, 621, 959
- Fruchter A. S., Stinebring D. R., Taylor J. H., 1988, *Nature*, 333, 237
- Harding A. K., Gaisser T. K., 1990, *ApJ*, 358, 561
- Hobbs G. B., Edwards R. T., Manchester R. N., 2006, *MNRAS*, 369, 655
- Iben Jr. I., Tutukov A. V., 1991, *ApJ*, 370, 615
- Janssen G. H., Stappers B. W., Kramer M., Nice D. J., Jessner A., Cognard I., Purver M. B., 2008, *A&A*, 490, 753
- Jeans J. H., 1924, *MNRAS*, 85, 2
- Keith M. J., Johnston S., Ray P. S., Ferrara E. C., Saz Parkinson P. M., Celik O., Belfiore A., Donato D., Cheung C. C., Abdo A. A., Camilo F., Freire P. C. C., Guillemot L., Harding A. K., Kramer M., Michelson P. F., Ransom S. M., Romani R. W., Smith D. A., Thompson D. J., Weltevrede P., Wood K. S., 2011, *ArXiv e-prints astro-ph/1102.0648*
- King A. R., Davies M. B., Beer M. E., 2003, *MNRAS*, 345, 678
- Kopeikin S. M., 1996, *ApJ*, 467, L93
- Lange C., Camilo F., Wex N., Kramer M., Backer D., Lyne A., Doroshenko O., 2001, *MNRAS*, 326, 274
- Lanza A. F., Rodonò M., 1999, *A&A*, 349, 887
- Lanza A. F., Rodonò M., 2001, *A&A*, 376, 165
- Lanza A. F., Rodono M., Rosner R., 1998, *MNRAS*, 296, 893
- Lazaridis K., Wex N., Jessner A., Kramer M., Stappers B. W., Janssen G. H., Desvignes G., Purver M. B., Cognard I., Theureau G., Lyne A. G., Jordan C. A., Zensus J. A., 2009, *MNRAS*, 400, 805
- Lorimer D. R., Kramer M., 2005, *Handbook of Pulsar Astronomy*. Cambridge University Press
- Nelemans G., Portegies Zwart S. F., Verbunt F., Yungelson L. R., 2001, *A&A*, 368, 939
- Nice D. J., Arzoumanian Z., Thorsett S. E., 2000, in Kramer M., Wex N., Wielebinski R., eds, *Pulsar Astronomy - 2000 and Beyond*, IAU Colloquium 177 Binary eclipsing millisecond pulsars: A decade of timing. Astronomical Society of the Pacific, San Francisco, pp 67–72
- Peters P. C., 1964, *Phys. Rev.*, 136, 1224
- Pfahl E., Rappaport S., Podsiadlowski P., 2002, *ApJ*, 573, 283
- Phinney E. S., 1992, *Philos. Trans. Roy. Soc. London A*, 341, 39
- Phinney E. S., Kulkarni S. R., 1994, *Ann. Rev. Astr. Ap.*, 32, 591
- Plyser E., Savonije G. J., 1988, *A&A*, 191, 57
- Ransom S. M., Ray P. S., Camilo F., Roberts M. S. E., Celik Ö., Wolff M. T., Cheung C. C., Kerr M., Pennucci T., DeCesar M. E., Cognard I., Lyne A. G., Stappers B. W., Freire P. C. C., Grove J. E., Abdo A. A., Desvignes G., Donato D., Ferrara E. C., Gehrels N., Guillemot L., Gwon C., Harding A. K., Johnston S., Keith M., Kramer M., Michelson P. F., Parent D., Saz Parkinson P. M., Romani R. W., Smith D. A., Theureau G., Thompson D. J., Weltevrede P., Wood K. S., Ziegler M., 2011, *ApJ*, 727, L16+
- Ritter H., 2008, *New Astron. Rev.*, 51, 869
- Savonije G. J., de Kool M., van den Heuvel E. P. J., 1986, *A&A*, 155, 51
- Shapiro S. L., Teukolsky S. A., 1983, *Black Holes, White Dwarfs and Neutron Stars. The Physics of Compact Objects*. Wiley–Interscience, New York
- Shklovskii I. S., 1970, *Sov. Astron.*, 13, 562
- Smarr L. L., Blandford R., 1976, *ApJ*, 207, 574
- Standish E. M., 1998, *JPL Planetary and Lunar Ephemerides, DE405/LE405*, Memo IOM 312.F-98-048. JPL, Pasadena
- Standish E. M., 2004, *A&A*, 417, 1165
- Stappers B. W., Bailes M., Lyne A. G., Camilo F., Manchester R. N., Sandhu J. S., Toscano M., Bell J. F., 2001a, *MNRAS*, 321, 576
- Stappers B. W., Bailes M., Lyne A. G., Manchester R. N., D’Amico N., Tauris T. M., Lorimer D. R., Johnston S., Sandhu J. S., 1996, *ApJ*, 465, L119
- Stappers B. W., Bailes M., Manchester R. N., Sandhu J. S., Toscano M., 1998, *ApJ*, 499, L183
- Stappers B. W., Bessell M. S., Bailes M., 1996, *ApJ*, 473, L119
- Stappers B. W., Gaensler B. M., Kaspi V. M., van der Klis M., Lewin W. H. G., 2003, *Science*, 299, 1372
- Stappers B. W., van Kerkwijk M. H., Bell J. F., Kulkarni S. R., 2001b, *ApJ*, 548, L183
- Stappers B. W., van Kerkwijk M. H., Lane B., Kulkarni S. R., 1999, *ApJ*, 510, L45
- Tauris T. M., Savonije G. J., 2001, in C. Kouveliotou, J. Ventura, & E. van den Heuvel ed., *The Neutron Star - Black Hole Connection Spin-Orbit Coupling in X-ray Binaries*. pp 337–+
- Tavani M., 1992, in van den Heuvel E. P. J., Rappaport S. A., eds, *X-ray Binaries and Recycled Pulsars Radiation-driven evolution of low-mass X-ray binaries and the formation of millisecond pulsars*. Kluwer, Dordrecht, pp 387–399
- van Kerkwijk M. H., Breton R. P., Kulkarni S. R., 2011, *ApJ*, 728, 95
- Wex N., 1998, *MNRAS*, 298, 67
- Wolszczan A., Frail D. A., 1992, *Nature*, 355, 145